CHAPTER 2 - SUPPLEMENTAL

HYPERSPECTRAL REMOTE SENSING OF WHITE MICA: A REVIEW OF IMAGING AND POINT-BASED SPECTROMETER STUDIES FOR MINERAL RESOURCES, WITH SPECTROMETER DESIGN CONSIDERATIONS.

A paper published in *Remote Sensing of Environment*

Reprinted with permission from Elsevier

John M Meyer[*1,2], Raymond F. Kokaly[2], Elizabeth Holley[1]

1 Chart References

Below are the references to be included with the '*S2_deposit_chart.xlsx*' included as a supplemental material.

1.1 References

Agus, A.J.L. (2011). Mapping white mica in milled porphyry copper pebbles using hyperspectral imagery: An exploratory study. In, Geo-Information Science and Earth 57. Enschede, The Netherlands: Universituy of Twente.

Ahmed, A.D., Hickey, K.A., & Barker, S.L. (2009). Tracing the distal hanging wall expression of Carlin-type hydrothermal. uids: a possible road-map to subsurface Au-deposits. Thermal and geochemical footprints of low-temperature sedimentary rock-hosted hydrothermal Au-systems: Identifying far-field vectors toward ore Year 2 Technical Summary. Vancouver, British Columbia: The University of British Columbia.

Bierwirth, P., Huston, D., & Blewett, R. (2002). Hyperspectral mapping of mineral assemblages associated with gold mineralization in the Central Pilbara, Western Australia. Economic Geology, 97 (4) 819-826. DOI: https://doi.org/10.2113/gsecongeo.97.4.819.

Brauhart, C., Huston, D., & Andrew, A. (2000). Oxygen isotope mapping in the Panorama VMS district, Pilbara Craton, Western Australia: applications to estimating temperatures of alteration and to exploration. Mineralium Deposita, 35 (8) 727-740. DOI: https://doi.org/10.1007/s001260050275.

---

[*]Primary researcher and author

[1]Department of Mining Engineering, Colorado School of Mines, Golden, CO, USA, 80401

[2] United States Geological Survey Box 25046 MS 973 Denver CO 80225-0046

Brill, B.A. (1988). Illite crystallinity, b0 and Si content of K-white mica as indicators of metamorphic conditions in low-grade metamorphic rocks at Cobar, New South Wales. Australian Journal of Earth Sciences, 35 (3) 295-302. DOI: https://doi.org/10.1080/08120098808729448.

Cline, J.S., Hofstra, A.H., Muntean, J.L., Tosdal, R.M., & Hickey, K.A. (2005). Carlin-type gold deposits in Nevada: Critical geologic characteristics and viable models. In J.W. Hedenquist, J.F.H. Thompson, R.J. Goldfarb, & J.P. Richards (Eds.), One Hundredth Anniversary Volume 484: Society of Economic Geologists. DOI: https://doi.org/10.5382/AV100.15.

Dalm, M., Buxton, M., & van Ruitenbeek, F. (2017). Discriminating ore and waste in a porphyry copper deposit using short-wavelength infrared (SWIR) hyperspectral imagery. Minerals engineering, 105 10-18. DOI: https://doi.org/10.1016/j.mineng.2016.12.013.

David, V. (2018). Cobar Deposits–Structural control. ASEG Extended Abstracts, 2018 (1) 1-10. DOI: http://dx.doi.org/10.1071/ASEG2018abT6_2G.

Duke, E.F. (1994). Near infrared spectra of muscovite, Tschermak substitution, and metamorphic reaction progress: Implications for remote sensing. Geology, 22 (7) 621-624. DOI: https://doi.org/10.1130/00917613(1994)022%3C0621:NISOMT%3E2.3.CO;2.

Duke, E.F., & Lewis, R.S. (2010). Near infrared spectra of white mica in the Belt Supergroup and implications for metamorphism. American Mineralogist, 95 (7) 908-920. DOI: http://dx.doi.org/10.2138/am.2010.3281.

Gaillard, N., Williams-Jones, A.E., Clark, J.R., Lypaczewski, P., Salvi, S., Perrouty, S., Piette-Lauzière, N., Guilmette, C., & Linnen, R.L. (2018). Mica composition as a vector to gold mineralization: Deciphering hydrothermal and metamorphic effects in the Malartic district, Quebec. Ore Geology Reviews, 95 789-820. DOI: https://doi.org/10.1016/j.oregeorev.2018.02.009.

Gifkins, C., & Allen, R. (2001). Textural and Chemical Characteristics of Diagenetic and Hydrothermal Alteration in Glassy Volcanic Rocks: Examples from the Mount Read Volcanics, Tasmania. Economic Geology, 96 (5) 973-1002. DOI: https://doi.org/10.2113/GSECONGEO.96.5.973.

Graham, G.E., Kokaly, R.F., Kelley, K.D., Hoefen, T.M., Johnson, M.R., & Hubbard, B.E. (2018). Application of imaging spectroscopy for mineral exploration in Alaska: A study over porphyry Cu deposits in the Eastern Alaska Range. Economic Geology, 113 (2) 489-510. DOI: https://doi.org/10.5382/econgeo.2018.4559.

Gregory, M.J., Lang, J.R., Gilbert, S., & Hoal, K.O. (2013). Geometallurgy of the Pebble porphyry copper-gold-molybdenum deposit, Alaska: Implications for gold distribution and paragenesis. Economic Geology, 108 (3) 463-482. DOI: http://dx.doi.org/10.2113/econgeo.108.3.463.

Guo, N., Thomas, C., Tang, J., & Tong, Q. (2019). Mapping white mica alteration associated with the Jiama porphyry-skarn Cu deposit, central Tibet using field SWIR spectrometry. Ore Geology Reviews, 108 11. DOI: https://doi.org/10.1016/j.oregeorev.2017.07.027.

Han, J., Chu, G., Chen, H., Hollings, P., Sun, S., & Chen, M. (2018). Hydrothermal alteration and short wavelength infrared (SWIR) characteristics of the Tongshankou porphyry-skarn Cu-Mo deposit, Yangtze craton, Eastern China. Ore Geology Reviews, 101 143-164. DOI: https://doi.org/10.1016/j.oregeorev.2018.07.018.

Harraden, C.L., Mcnulty, B.A., Gregory, M.J., & Lang, J.R. (2013). Shortwave infrared spectral analysis of hydrothermal alteration associated with the Pebble porphyry copper-gold-molybdenum deposit, Iliamna, Alaska. Economic Geology, 108 (3) 483-494. DOI: https://doi.org/10.2113/econgeo.108.3.483.

Herrmann, W., Blake, M., Doyle, M., Huston, D., Kamprad, J., Merry, N., & Pontual, S. (2001). Short wavelength infrared (SWIR) spectral analysis of hydrothermal alteration zones associated with base metal sulfide deposits at Rosebery and Western Tharsis, Tasmania, and Highway-Reward, Queensland. Economic Geology, 96 (5) 939-954. DOI: http://dx.doi.org/10.2113/96.5.939.

Hewson, R., Cudahy, T., Mizuhiko, S., Ueda, K., & Mauger, A. (2005). Seamless geological map generation using ASTER in the Broken Hill-Curnamona province of Australia. Remote Sensing of Environment, 99 (1-2) 159-172.DOI: https://doi.org/10.1016/j.rse.2005.04.025.

Hitzman, M.W., Oreskes, N., & Einaudi, M.T. (1992). Geological characteristics and tectonic setting of proterozoic iron oxide (Cu-U-Au-REE) deposits. Precambrian Research, 58 (1-4) 241-287. DOI: https://doi.org/10.1016/0301-9268(92)90121-4.

Huang, J., Chen, H., Han, J., Deng, X., Lu, W., & Zhu, R. (2018). Alteration zonation and short wavelength infrared (SWIR) characteristics of the Honghai VMS Cu-Zn deposit, Eastern Tianshan, NW China. Ore Geology Reviews, 100 263-279. DOI: http://dx.doi.org/10.1016/j.oregeorev.2017.02.037.

Huston, D.L., Blewett, R.S., Keillor, B., Standing, J., Smithies, R.H., Marshall, A., Mernagh, T.P., & Kamprad, J. (2002). Lode gold and epithermal deposits of the Mallina basin, North Pilbara terrain, Western Australia. Economic Geology, 97 (4) 801-818. DOI: http://dx.doi.org/10.2113/gsecongeo.97.4.801.

Huston, D.L., Kamprad, J.L., & Brauhart, C. (1997). Preliminary Results of PIMA Analyses of the Alteration Zone Underlying the Sulphur Springs Desposit, Panorama District, Pilbara Block, Western Australia. Canberra, Australia: Australian Geological Survey Organisation.

Jones, S., Herrmann, W., & Gemmell, J.B. (2005). Short wavelength infrared spectral characteristics of the HW horizon: Implications for exploration in the Myra Falls volcanic-hosted massive sulfide camp, Vancouver Island, British Columbia, Canada. Economic Geology, 100 (2) 273-294. DOI: https://doi.org/10.2113/gsecongeo.100.2.273.

Kokaly, R.F., Graham, G.E., Hoefen, T.M., Kelley, K.D., Johnson, M.R., Hubbard, B.E., Buchhorn, M., & Prakash, A. (2017). Multiscale Hyperspectral Imaging of the Orange Hill Porphyry Copper Deposit. In, Proceedings of Exploration 17: Sixth Decennial International Conference on Mineral Exploration 923-943: Decennial Mineral Exploration Conferences.

Komninou, A., & Sverjensky, D. (1995). Hydrothermal alteration and the chemistry of ore-forming fluids in an unconformity-type uranium deposit. Geochimica et Cosmochimica Acta, 59 (13) 2709-2723. DOI: https://doi.org/10.1016/0016-7037(95)00167-X.

Kruse, F.A., Bedell, R.L., Taranik, J.V., Peppin, W.A., Weatherbee, O., & Calvin, W.M. (2012). Mapping alteration minerals at prospect, outcrop and drill core scales using imaging spectrometry. International Journal of Remote Sensing, 33 (6) 1780-1798. DOI: https://dx.doi.org/10.1080%2F01431161.2011.600350.

Kruse, F.A., & Hauff, P.L. (1991). Identification of illite polytype zoning in disseminated gold deposits using reflectance spectroscopy and X-ray diffraction-potential for mapping with imaging spectrometers. IEEE Transactions on Geoscience and Remote Sensing, 29 (1) 101-104. DOI: http://dx.doi.org/10.1109/36.103298.

Laakso, K., Peter, J., Rivard, B., & White, H. (2016). Short-wave infrared spectral and geochemical characteristics of hydrothermal alteration at the Archean Izok Lake Zn-Cu-Pb-Ag volcanogenic massive sulfide deposit, Nunavut, Canada: Application in exploration target vectoring. Economic Geology, 111 (5) 1223-1239. DOI: http://dx.doi.org/10.2113/econgeo.111.5.1223.

Lang, J.R., Gregory, M.J., Rebagliati, C.M., Payne, J.G., Oliver, J.L., & Roberts, K. (2013). Geology and magmatic-hydrothermal evolution of the giant Pebble porphyry copper-gold-molybdenum deposit, southwest Alaska. Economic Geology, 108 (3) 437-462. DOI: http://dx.doi.org/10.2113/econgeo.108.3.437.

Lawrie, K., & Hinman, M. (1998). Cobar-style polymetallic Au-Cu-Ag-Pb-Zn deposits. AGSO Journal of Australian Geology and Geophysics, 17 (4) 169-187.

Lypaczewski, P., Rivard, B., Gaillard, N., Perrouty, S., Piette-Lauzière, N., Bérubé, C.L., & Linnen, R.L. (2019). Using hyperspectral imaging to vector towards mineralization at the Canadian Malartic gold deposit, Québec, Canada. Ore Geology Reviews, 111 15. DOI: https://doi.org/10.1016/j.oregeorev.2019.102945.

Mateer, M.A. (2010). Ammonium illite at the Jerritt Canyon district and Goldstrike property, Nevada: Its spatial distribution and significance in the exploration of Carlin-type deposits. In, Geology and Geophysics 214: University of Wyoming.

Mathieu, M., Roy, R., Launeau, P., Cathelineau, M., & Quirt, D. (2017). Alteration mapping on drill cores using a HySpex SWIR-320m hyperspectral camera: Application to the exploration of an unconformity-related uranium deposit (Saskatchewan, Canada). Journal of Geochemical Exploration, 172 71-88. DOI: https://doi.org/10.1016/j.gexplo.2016.09.008.

Post, J.L., & Noble, P.N. (1993). The near-infrared combination band frequencies of dioctahedral smectites, micas, and illites. Clays and Clay Minerals, 41 639-640. DOI: https://doi.org/10.1346/CCMN.1993.0410601.

Robson, D., Cocks, P., & Taylor, G. (2003). Calibration, processing and interpretation of hyperspectral data over the Broken Hill region. In, ASEG 16th Geophysical Conference and Exhibition. Adelaide, Australia: ASEG Extended Abstracts. DOI: https://doi.org/10.1071/ASEG2003ab149.

Squire, R.J., Herrmann, W., Pape, D., & Chalmers, D.I. (2007). Evolution of the Peak Hill high-sulfidation epithermal Au–Cu deposit, Eastern Australia. Mineralium Deposita, 42 (5) 489-503. DOI: http://dx.doi.org/10.1007/s00126-006-0076-4.

Sun, Y., Seccombe, P.K., & Yang, K. (2001). Application of short-wave infrared spectroscopy to define alteration zones associated with the Elura zinc–lead–silver deposit, NSW, Australia. Journal of Geochemical Exploration, 73 (1) 11-26. DOI: https://doi.org/10.1016/S0375-6742(01)00167-4.

Swayze, G.A., Clark, R.N., Goetz, A.F., Livo, K.E., Breit, G.N., Kruse, F.A., Sutley, S.J., Snee, L.W., Lowers, H.A., Post, J.L., Stoffregen, R.E., & Ashley, R.P. (2014). Mapping advanced argillic alteration at Cuprite, Nevada, using imaging spectroscopy. Economic Geology, 109 (5) 1179-1222. DOI: http://dx.doi.org/10.2113/econgeo.109.5.1179.

Tappert, M., Rivard, B., Giles, D., Tappert, R., & Mauger, A. (2011). Automated drill core logging using visible and near-infrared reflectance spectroscopy: a case study from the Olympic Dam IOCG deposit, South Australia. Economic Geology, 106 (2) 289-296. DOI: http://dx.doi.org/10.2113/econgeo.106.2.289.

Tappert, M.C., Rivard, B., Giles, D., Tappert, R., & Mauger, A. (2013). The mineral chemistry, near-infrared, and mid-infrared reflectance spectroscopy of phengite from the Olympic Dam IOCG deposit, South Australia. Ore Geology Reviews, 53 26-38. DOI: http://dx.doi.org/10.1016/j.oregeorev.2012.12.006.

Uribe-Mogollon, C., & Maher, K. (2020). White Mica Geochemistry: Discriminating Between Barren and Mineralized Porphyry Systems. Economic Geology, 115 (2) 325-354. DOI: http://dx.doi.org/10.5382/econgeo.4706.

van der Meer, F., Kopačková, V., Koucká, L., van der Werff, H.M., van Ruitenbeek, F.J., & Bakker, W.H. (2018). Wavelength feature mapping as a proxy to mineral chemistry for investigating geologic systems: An example from the Rodalquilar epithermal system. International Journal of Applied Earth Observation and Geoinformation, 64 237-248. DOI: https://doi.org/10.1016/j.jag.2017.09.008.

van Ruitenbeek, F.J., Cudahy, T., Hale, M., & van der Meer, F.D. (2005). Tracing fluid pathways in fossil hydrothermal systems with near-infrared spectroscopy. Geology, 33 (7) 597-600. DOI: https://doi.org/10.1130/G21375.1.

van Ruitenbeek, F.J., Debba, P., van der Meer, F.D., Cudahy, T., van der Meijde, M., & Hale, M. (2006). Mapping white micas and their absorption wavelengths using hyperspectral band ratios. Remote Sensing of Environment, 102 (3-4) 211-222. DOI: https://doi.org/10.1016/j.rse.2006.02.012.

Yang, K., Huntington, J., Boardman, J., & Mason, P. (1999). Mapping hydrothermal alteration in the Comstock mining district, Nevada, using simulated satellite-borne hyperspectral data. Australian Journal of Earth Sciences, 46 (6) 915-922. DOI: http://dx.doi.org/10.1046/j.1440-0952.1999.00754.x.

Yang, K., Huntington, J., Gemmell, J., & Scott, K. (2011). Variations in composition and abundance of white mica in the hydrothermal alteration system at Hellyer, Tasmania, as

revealed by infrared reflectance spectroscopy. Journal of Geochemical Exploration, 108 (2) 143-156. DOI: http://dx.doi.org/10.1016/j.gexplo.2011.01.001.